

The use of carboxylic acid derivatives as drugs

This application is a 371 of PCT/EP95/01099 filed on March 29, 1995.

The present invention relates to the use of certain carboxylic acid derivatives as drugs.

Endothelin is a peptide which is composed of 21 amino acids and which is synthesized and released by vascular endothelium. Endothelin exists in three isoforms, ET-1, ET-2 and ET-3. "Endothelin" or "ET" hereinafter means one or all isoforms of endothelin. Endothelin is a potent vasoconstrictor and has a potent effect on vascular tone. It is known that this vasoconstriction is caused by binding of endothelin to its receptor (Nature 332 (1988) 411-415; FEBS Letters 231 (1988) 440-444, and Biochem. Biophys. Res. Commun. 154 (1988) 868-875).


Increased or abnormal release of endothelin causes persistent vasoconstriction in peripheral, renal and cerebral vessels, which may lead to pathological states. It is reported in the literature that elevated plasma endothelin levels are found in patients with hypertension, acute myocardial infarct, pulmonary hypertension, Raynaud's syndrome or atherosclerosis and in the airways of asthmatics (Japan J. Hypertension 12 (1989) 79, J. Vascular Med. Biology 2 (1990) 207, J. Am. Med. Association 264 (1990) 2868).

Accordingly, substances which specifically inhibit the binding of endothelin to the receptor should also antagonize the various physiological effects of endothelin mentioned above and therefore be valuable drugs.

We have found that certain carboxylic acid derivatives are good inhibitors of endothelin receptors.

The invention relates to the use of carboxylic acid derivatives with the formula I which is described hereinafter for the production of drugs, in particular for the production of inhibitors of endothelin receptors.

Carboxylic acid derivatives of the general formula I



where R is formyl, CO₂H or a radical which can be hydrolyzed to COOH, and the remaining substituents have the following meanings:

- 5 R² is halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;
- 10 X is nitrogen or CR¹⁴ where R¹⁴ is hydrogen or, together with R³, forms a 3- or 4-membered alkylene or alkenylene chain in which, in each case, one methylene group is replaced by oxygen;
- 15 R³ is halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio or R³ is linked to R¹⁴ as indicated above to form a 5- or 6-membered ring;
- 20 R⁴ is C₁-C₁₀-alkyl which can carry from one to five halogen atoms and/or one of the following radicals: C₁-C₄-alkoxy, C₁-C₄-alkylthio, cyano, C₁-C₈-alkylcarbonyl, C₁-C₈-alkoxycarbonyl, phenyl, phenoxy or phenylcarbonyl, where the phenyl radicals in turn can carry from one to five halogen atoms and/or from one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;
- 25 C₁-C₁₀-alkyl which can carry from one to five halogen atoms and carries one of the following radicals: a five-membered heteroaromatic ring which contains from one to three nitrogen atoms and/or one sulfur or oxygen atom and which can carry from one to four halogen atoms and/or one or two of the fol-
- 30 lowing radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio and/or phenyl;
- 35 C₃-C₁₂-cycloalkyl or C₃-C₁₂-cycloalkenyl, each of which can contain one oxygen or sulfur atom and can carry from one to five halogen atoms and/or one of the following radicals: C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, cyano, C₁-C₈-alkylcarbonyl, C₁-C₈-alkoxycarbonyl, phenyl, phenoxy or phenylcarbonyl, where the phenyl radicals in turn can carry from one to five halogen atoms and/or from one to three of the
- 40 following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;
- 45 C₃-C₆-alkenyl or C₃-C₆-alkynyl, each of which can carry from one to five halogen atoms and/or one of the following radicals: C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, cyano, C₁-C₈-alkylcarbonyl, C₁-C₈-alkoxycarbonyl, phenyl, phenoxy or phenylcarbonyl, where the phenyl radicals in turn can carry

from one to five halogen atoms and/or from one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;

- 5 a five- or six-membered heteroaromatic ring which contains from one to three nitrogen atoms and/or one sulfur or oxygen atom and which can carry from one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, phenyl, phenoxy or phenylcarbonyl, where the phenyl radicals in turn can carry from one to five halogen atoms and/or from one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;
- 10
- 15 phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, amino, C₁-C₄-alkylamino or C₁-C₄-dialkylamino;
- 20

- R⁴ and R⁵ form, together with the adjacent carbon atom, a 3- to 8-membered ring which can contain one oxygen or sulfur atom and can carry from one to three of the following radicals: C₁-C₄-alkyl, halogen, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio [sic];
- 25

- R⁵ is hydrogen, C₁-C₄-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₈-cycloalkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxyalkyl, C₁-C₄-alkylthioalkyl, phenyl or R⁵ is linked to R⁴ as indicated above to form a 3- to 8-membered ring;
- 30

- R⁶ is C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or C₃-C₈-cycloalkyl, it being possible for each of these radicals to be substituted one or more times by: halogen, nitro, cyano, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxy-carbonyl, C₁-C₄-alkylamino, di-C₁-C₄-alkylamino, phenyl, phenoxy or phenyl which is substituted one or more times, eg. from one to three times, by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio;
- 35
- 40

- phenyl or naphthyl, each of which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy,
- 45


C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino or C₁-C₄-dialkylamino;

a five- or six-membered heteroaromatic ring which contains from one to three nitrogen atoms and/or one sulfur or oxygen atom and which can carry from one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, phenyl, phenoxy or phenylcarbonyl, where the phenyl radicals in turn can carry from one to five halogen atoms and/or from one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio;

Y is sulfur or oxygen or a single bond;

Z is sulfur or oxygen.


The compounds according to the invention are prepared starting from the epoxides IV which are obtained in a conventional manner, eg. as described in J. March, Advanced Organic Chemistry, 2nd ed., 1983, p. 862 and p. 750, from the aldehydes or ketones II or the olefins III:



Carboxylic acid derivatives of the general formula VI can be prepared by reacting the epoxides of the general formula IV (eg. with R = COOR¹⁰) with alcohols or thiols of the general formula V where R⁶ and Z have the meanings classified in claim 1.

6

5



10


where R^{15} is halogen or $\text{R}^{16}-\text{SO}_2-$, where R^{16} can be C_1-C_4 -alkyl, C_1-C_4 -haloalkyl or phenyl. The reaction preferably takes place in one of the abovementioned inert diluents with the addition of a suitable base, ie. a base which deprotonates the intermediate VI, 15 at from room temperature to the boiling point of the solvent.

The base which can be used is an alkali metal or alkaline earth metal hydride such as sodium hydride, potassium hydride or calcium hydride or a carbonate such as an alkali metal carbonate, 20 eg. sodium or potassium carbonate, an alkali metal or alkaline earth metal hydroxide such as sodium or potassium hydroxide, an organometallic compound such as butyllithium or an alkali metal amide such as lithium diisopropylamide.

25 The compounds according to the invention where Y is sulfur and the remaining substituents have the meanings indicated for the general formula I can be prepared, for example, by reacting carboxylic acid derivatives of the general formula VIII, which can be obtained in a conventional manner from compounds of the 30 general formula VI and in which the substituents have the above-mentioned meanings, with compounds of the general formula IX where R^2 , R^3 and X have the meanings indicated for the general formula I.

35

40



45

The reaction preferably takes place in one of the abovementioned inert diluents with the addition of a suitable base, ie. a base which deprotonates the intermediate IX, at from room temperature to the boiling point of the solvent.


5

Besides the abovementioned bases it is also possible to use organic bases such as tertiary amines, eg. triethylamine, pyridine, imidazole or diazabicycloandecene.

- 10 Compounds of the formula I can also be prepared by starting from the corresponding carboxylic acids, ie. compounds of the formula I where R^1 is hydroxyl, and initially converting these in a conventional way into an activated form, such as a halide, an anhydride or imidazolidine, and then reacting the latter with an
15 appropriate hydroxyl compound HOR^{10} . This reaction can be carried out in the conventional solvents and often requires addition of a base, in which case those mentioned above are suitable. These two steps can also be simplified, for example, by allowing the carboxylic acid to act on the hydroxyl compound in the presence
20 of a dehydrating agent such as a carbodiimide.

Compounds of the formula I can also be prepared by starting from salts of the appropriate carboxylic acids, ie. from compounds of the formula I where R is COR^1 and R^1 is OM where M can be an
25 alkali metal cation or the equivalent of an alkaline earth metal cation. These salts can be reacted with many compounds of the formula R^1-A where A is a conventional nucleofugic leaving group, for example halogen such as chlorine, bromine, iodine or aryl- or alkylsulfonyl which is unsubstituted or substituted by halogen,
30 alkyl or haloalkyl, such as toluenesulfonyl and methylsulfonyl, or another equivalent leaving group. Compounds of the formula R^1-A with a reactive substituent A are known or can easily be obtained using general expert knowledge. This reaction can be carried out in the conventional solvents, advantageously with the addition of
35 a base, those mentioned above being suitable.

The radical R in formula I can vary widely. R is, for example,

T80X
40

where R^1 has the following meanings:

45

a) hydrogen;

b) succinimidyloxy;

c) a 5-membered heteroaromatic ring linked via a nitrogen atom, such as pyrrolyl, pyrazolyl, imidazolyl and triazolyl, which can carry one or two halogen atoms, especially fluorine and chlorine and/or one or two of the following radicals:

C₁-C₄-alkyl such as methyl, ethyl, 1-propyl, 2-propyl, 2-methyl-2-propyl, 2-methyl-1-propyl, 1-butyl, 2-butyl;


C₁-C₄-haloalkyl, in particular C₁-C₂-haloalkyl such as fluoro-methyl, difluoromethyl, trifluoromethyl, chlorodifluoro-methyl, dichlorofluoromethyl, trichloromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and pentafluoroethyl;

C₁-C₄-haloalkoxy, in particular C₁-C₂-haloalkoxy such as di-fluoromethoxy, trifluoromethoxy, chlorodifluoromethoxy, 1-fluoroethoxy, 2-fluoroethoxy, 2,2-difluoroethoxy, 1,1,2,2-tetrafluoroethoxy, 2,2,2-trifluoroethoxy, 2-chloro-1,1,2-trifluoroethoxy and pentafluoroethoxy, especially trifluoromethoxy;

C₁-C₄-alkoxy such as methoxy, ethoxy, propoxy, 1-methylethoxy, butoxy, 1-methylpropoxy, 2-methylpropoxy, 1,1-dimethylethoxy, especially methoxy, ethoxy, 1-methylethoxy;

C₁-C₄-alkylthio such as methylthio, ethylthio, propylthio, 1-methylethylthio, butylthio, 1-methylpropylthio, 2-methyl-propylthio, 1,1-dimethylethylthio, especially methylthio and ethylthio;

d)



where m is 0 or 1 and R⁷ and R⁸, which can be identical or different, have the following meanings:

hydrogen

T90X

C₁-C₈-alkyl, especially C₁-C₄-alkyl as mentioned above;

C₃-C₆-alkenyl such as 2-propenyl, 2-butenyl, 3-butenyl,
 1-methyl-2-propenyl, 2-methyl-2-propenyl, 2-pentenyl,
 5 3-pentenyl, 4-pentenyl, 1-methyl-2-butenyl,
 2-methyl-2-butenyl, 3-methyl-2-butenyl, 1-methyl-3-butenyl,
 2-methyl-3-butenyl, 3-methyl-3-butenyl, 1,1-dimethyl-2-
 propenyl, 1,2-dimethyl-2-propenyl, 1-ethyl-2-propenyl,
 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 1-methyl-
 10 2-pentenyl, 2-methyl-2-pentenyl, 3-methyl-2-pentenyl,
 4-methyl-2-pentenyl, 3-methyl-3-pentenyl, 4-methyl-3-
 pentenyl, 1-methyl-4-pentenyl, 2-methyl-4-pentenyl,
 3-methyl-4-pentenyl, 4-methyl-4-pentenyl,
 1,1-dimethyl-2-butenyl, 1,1-dimethyl-3-butenyl,
 15 1,2-dimethyl-2-butenyl, 1,2-dimethyl-3-butenyl,
 1,3-dimethyl-2-butenyl, 1,3-dimethyl-3-butenyl,
 2,2-dimethyl-3-butenyl, 2,3-dimethyl-2-butenyl,
 2,3-dimethyl-3-butenyl, 1-ethyl-2-butenyl, 1-ethyl-3-butenyl,
 2-ethyl-2-butenyl, 2-ethyl-3-butenyl, 1,1,2-trimethyl-2-pro-
 20 penyl, 1-ethyl-1-methyl-2-propenyl and 1-ethyl-2-methyl-
 2-propenyl, especially 2-propenyl, 2-butenyl, 3-methyl-
 2-butenyl and 3-methyl-2-pentenyl;

C₃-C₆-alkynyl such as 2-propynyl, 2-butyngyl, 3-butyngyl,
 25 1-methyl-2-propynyl, 2-pentyngyl, 3-pentyngyl, 4-pentyngyl,
 1-methyl-3-butyngyl, 2-methyl-3-butyngyl, 1-methyl-2-butyngyl,
 1,1-dimethyl-2-propynyl, 1-ethyl-2-propynyl, 2-hexynyl,
 3-hexynyl, 4-hexynyl, 5-hexynyl, 1-methyl-2-pentyngyl,
 1-methyl-3-pentyngyl, 1-methyl-4-pentyngyl, 2-methyl-
 30 3-pentyngyl, 2-methyl-4-pentyngyl, 3-methyl-4-pentyngyl,
 4-methyl-2-pentyngyl, 1,1-dimethyl-2-butyngyl,
 1,1-dimethyl-3-butyngyl, 1,2-dimethyl-3-butyngyl,
 2,2-dimethyl-3-butyngyl, 1-ethyl-2-butyngyl, 1-ethyl-3-butyngyl,
 2-ethyl-3-butyngyl and 1-ethyl-1-methyl-2-propynyl, preferably
 35 2-propynyl, 2-butyngyl, 1-methyl-2-propynyl and 1-methyl-
 2-butyngyl, especially 2-propynyl

C₃-C₈-cycloalkyl, such as cyclopropyl, cyclobutyl, cyclo-
 pentyl, cyclohexyl, cycloheptyl and cyclooctyl, it being
 40 possible for these alkyl, cycloalkyl, alkenyl and alkynyl
 groups in each case to carry from one to five halogen atoms,
 especially fluorine or chlorine, and/or one or two of the
 following groups:

45 C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy as
 mentioned above, C₃-C₆-alkenyloxy, C₃-C₆-alkenylthio,
 C₃-C₆-alkynyloxy, C₃-C₆-alkynylthio, where the alkenyl and

alkynyl moieties present in these radicals preferably have the abovementioned meanings;

5 C₁-C₄-alkylcarbonyl such as, in particular, methylcarbonyl, ethylcarbonyl, propylcarbonyl, 1-methylethylcarbonyl, butylcarbonyl, 1-methylpropylcarbonyl, 2-methylpropylcarbonyl, 1,1-dimethylethylcarbonyl;

10 C₁-C₄-alkoxycarbonyl such as methoxycarbonyl, ethoxycarbonyl, propyloxycarbonyl, 1-methylethoxycarbonyl, butyloxycarbonyl, 1-methylpropyloxycarbonyl, 2-methylpropyloxycarbonyl, 1,1-dimethylethoxycarbonyl;

15 C₃-C₆-alkenylcarbonyl, C₃-C₆-alkynylcarbonyl, C₃-C₆-alkenyloxy-carbonyl and C₃-C₆-alkynyloxycarbonyl, where the alkenyl and alkynyl radicals are preferably defined as detailed above;

20 phenyl which is unsubstituted or substituted one or more times, eg. from once to three times, by halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio, such as 2-fluorophenyl, 3-chlorophenyl, 4-bromophenyl, 2-methylphenyl, 3-nitrophenyl, 4-cyanophenyl, 2-trifluoromethylphenyl, 3-methoxyphenyl, 4-trifluoroethoxyphenyl, 2-methylthiophenyl, 2,4-dichloro-phenyl, 2-methoxy-3-methylphenyl, 2,4-dimethoxyphenyl, 25 2-nitro-5-cyanophenyl, 2,6-difluorophenyl;

30 di-C₁-C₄-alkylamino such as, in particular, dimethylamino, di-propylamino, N-propyl-N-methylamino, N-propyl-N-ethylamino, diisopropylamino, N-isopropyl-N-methylamino, N-isopropyl-N-ethylamino, N-isopropyl-N-propylamino;


35 R⁷ and R⁸ are also phenyl which can be substituted by one or more, eg. from one to three, of the following radicals: halogen, nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy or C₁-C₄-alkylthio as mentioned above in particular;

40 or R⁷ and R⁸ together form a C₄-C₇-alkylene chain which is closed to form a ring and is unsubstituted or substituted, eg. by C₁-C₄-alkyl, and can contain a hetero atom selected from the group comprising oxygen, sulfur or nitrogen, such as -(CH₂)₄-, -(CH₂)₅-, -(CH₂)₆-, -(CH₂)₇-, -(CH₂)₂-O-(CH₂)₂-, -CH₂-S-(CH₂)₃-, -(CH₂)₂-O-(CH₂)₃-, -NH-(CH₂)₃-, -CH₂-NH-(CH₂)₂-, 45 -CH₂-CH=CH-CH₂-, -CH=CH-(CH₂)₃-;

e)

T120X

5



where k is 0, 1 or 2, p is 1, 2, 3 or 4, and R⁹ is

10 C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or unsubstituted or substituted phenyl as mentioned above in particular.

f) R¹ is also OR¹⁰ where R¹⁰ is:

15 hydrogen, the cation of an alkali metal such as lithium, sodium, potassium or the cation of an alkaline earth metal such as calcium, magnesium and barium or an environmentally compatible organic ammonium ion such as tertiary C₁-C₄-alkylammonium or the ammonium ion;

20

C₃-C₈-cycloalkyl as mentioned above, which can carry from one to three C₁-C₄-alkyl groups;

25

C₁-C₈-alkyl such as, in particular, methyl, ethyl, propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethylbutyl, 2-ethylbutyl, 1-ethyl-2-methylpropyl, which can carry from one to five halogen atoms, in particular fluorine and chlorine, and/or

30

35

C₁-C₄-alkoxy, C₁-C₄-alkylthio, cyano, C₁-C₄-alkylcarbonyl, C₃-C₈-cycloalkyl, C₁-C₄-alkoxycarbonyl, phenyl, phenoxy or phenylcarbonyl, where the aromatic radicals can in turn each carry from one to five halogen atoms and/or from one to three of the following radicals: nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio, in particular as mentioned above;

40

45

C₁-C₈-alkyl as mentioned above, which can carry from one to five halogen atoms, in particular fluorine and/or chlorine, and carries one of the following radicals: a 5-membered

heteroaromatic ring containing from one to three nitrogen atoms, or a 5-membered heteroaromatic ring containing one nitrogen atom and one oxygen or sulfur atom, which can carry from one to four halogen atoms and/or one or two of the following radicals:

nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, phenyl, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio. The following may be particularly mentioned: 1-pyrazolyl, 3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl, 1-benzotriazolyl, 3-isopropyl-5-isoxazolyl, 3-methyl-5-isoxazolyl, 2-oxazolyl, 2-thiazolyl, 2-imidazolyl, 3-ethyl-5-isoxazolyl, 3-phenyl-5-isoxazolyl, 3-tert-butyl-5-isoxazolyl;


C₂-C₆-alkyl which carries one of the following radicals in position 2: C₁-C₄-alkoxyimino, C₃-C₆-alkynyloxyimino, C₃-C₆-haloalkenyloxyimino or benzyloxyimino;

C₃-C₆-alkenyl or C₃-C₆-alkynyl, where these groups in turn can carry from one to five halogen atoms;

R¹⁰ is also a phenyl which can carry from one to five halogen atoms and/or from one to three of the following radicals: nitro, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio, in particular as mentioned above;

a 5-membered heteroaromatic ring which is linked via a nitrogen atom, contains from one to three nitrogen atoms and can carry one or two halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, phenyl, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio. The following may be particularly mentioned: 1-pyrazolyl, 3-methyl-1-pyrazolyl, 4-methyl-1-pyrazolyl, 3,5-dimethyl-1-pyrazolyl, 3-phenyl-1-pyrazolyl, 4-phenyl-1-pyrazolyl, 4-chloro-1-pyrazolyl, 4-bromo-1-pyrazolyl, 1-imidazolyl, 1-benzimidazolyl, 1,2,4-triazol-1-yl, 3-methyl-1,2,4-triazol-1-yl, 5-methyl-1,2,4-triazol-1-yl, 1-benzotriazolyl, 3,4-dichloro-1-imidazolyl;

R¹⁰ is also a group



5

where R^{11} and R^{12} , which can be identical or different, are:

10

$\text{C}_1\text{--C}_8\text{--alkyl}$, $\text{C}_3\text{--C}_6\text{--alkenyl}$, $\text{C}_3\text{--C}_6\text{--alkynyl}$, $\text{C}_3\text{--C}_8\text{--cycloalkyl}$, it being possible for these radicals to carry a $\text{C}_1\text{--C}_4\text{--alkoxy}$, $\text{C}_1\text{--C}_4\text{--alkylthio}$ and/or a substituted or unsubstituted phenyl radical, in particular as mentioned above;

15


phenyl, which can be substituted by one or more, eg. from one to three, of the following radicals: halogen, nitro, cyano, $\text{C}_1\text{--C}_4\text{--alkyl}$, $\text{C}_1\text{--C}_4\text{--haloalkyl}$, $\text{C}_1\text{--C}_4\text{--alkoxy}$, $\text{C}_1\text{--C}_4\text{--haloalkoxy}$ or $\text{C}_1\text{--C}_4\text{--alkylthio}$, where these radicals correspond in particular to those mentioned above;

20

or R^{11} and R^{12} together form a $\text{C}_3\text{--C}_{12}\text{--alkylene}$ chain which can carry from one to three $\text{C}_1\text{--C}_4\text{--alkyl}$ groups and contain a hetero atom from the group comprising oxygen, sulfur and nitrogen, in particular as mentioned for R^7 and R^8 .

g) R^1 is also

25



30

where R^{13} is:

35

$\text{C}_1\text{--C}_4\text{--alkyl}$, $\text{C}_3\text{--C}_6\text{--alkenyl}$, $\text{C}_3\text{--C}_6\text{--alkynyl}$, $\text{C}_3\text{--C}_8\text{--cycloalkyl}$, in particular as mentioned above, it being possible for these radicals to carry a $\text{C}_1\text{--C}_4\text{--alkoxy}$, $\text{C}_1\text{--C}_4\text{--alkylthio}$ and/or a phenyl radical as mentioned above;

40

phenyl which is unsubstituted or substituted, in particular as mentioned above.

With a view to the biological effect, preferred carboxylic acid derivatives of the general formula I are those in which the substituents have the following meanings:

45

- 5 R^2 the C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy, C_1-C_4 -halo-alkoxy, C_1-C_4 -alkylthio groups and halogen atoms mentioned specifically for R^1 , in particular chlorine, methyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy, particularly preferably methoxy;
- X nitrogen or CR^{14} where
- 10 R^{14} is hydrogen or forms together with R^3 a 4- or 5-membered alkylene or alkenylene chain in which, in each case, one methylene group is replaced by oxygen, such as $-CH_2-CH_2-O-$, $-CH=CH-O-$, $-CH_2-CH_2-CH_2-O-$, $-CH=CH-CH_2O-$, in particular hydrogen and $-CH_2-CH_2-O-$;
- 15 R^3 the C_1-C_4 -alkyl, C_1-C_4 -haloalkyl, C_1-C_4 -alkoxy, C_1-C_4 -halo-alkoxy, C_1-C_4 -alkylthio groups and halogen atoms mentioned for R^1 , in particular chlorine, methyl, methoxy, ethoxy, difluoromethoxy, trifluoromethoxy or is linked to R^{14} as mentioned above to form a 5- or 6-membered ring, R^3 is particularly preferably methoxy;
- 20 R^4 C_1-C_{10} -alkyl as specifically mentioned for R^1 , which can carry from one to five halogen atoms such as fluorine, chlorine, bromine, iodine, in particular fluorine and chlorine, and/or
- 25 one of the following radicals: alkoxy, alkylthio, cyano, alkylcarbonyl, alkoxycarbonyl, phenyl, phenoxy, phenyl-carbonyl as mentioned in general and in particular for R^1 ;
- 30 C_1-C_{10} -alkyl as mentioned above, which can carry from one to five halogen atoms as mentioned above, in particular fluorine and chlorine, and carries a 5-membered heteroaromatic ring which is unsubstituted or substituted, as mentioned above for R^1 ;
- 35 C_3-C_{12} -cycloalkyl, in particular C_3-C_7 -cycloalkyl, or C_3-C_{12} -cycloalkenyl, in particular C_4-C_7 -cycloalkenyl, it being possible for one methylene group in the saturated or unsaturated ring to be replaced by an oxygen or sulfur atom, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, cyclopropenyl, dihydrofuranyl, dihydrothienyl, dihydropyranyl, dihydrothiopyranyl,
- 40 where the cycloalkyl and cycloalkenyl radicals can be substituted by from one to five halogen atoms as mentioned above, especially fluorine or chlorine, and/or one of the following
- 45 radicals: C_1-C_4 -alkyl, C_1-C_4 -alkoxy, C_1-C_4 -alkylthio, cyano, C_1-C_8 -alkylcarbonyl, C_1-C_8 -alkoxycarbonyl, phenyl, phenoxy,

phenylcarbonyl as mentioned above in general and in particular;

5 C₃-C₆-alkenyl or C₃-C₆-alkynyl as mentioned for R¹, which can carry from one to five halogen atoms as mentioned above, in particular fluorine and chlorine, and/or one of the following radicals:

10 C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, cyano, C₁-C₈-alkylcarbonyl, C₁-C₈-alkoxycarbonyl, phenyl, phenoxy, phenylcarbonyl as mentioned above in general and in particular;

15 5- or 6-membered hetaryl such as furyl, thienyl, pyrrol, pyrazolyl, imidazolyl, triazolyl, isoxazolyl, oxazolyl, isothiazolyl, thiazolyl, thiadiazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, for example 2-furanyl, 3-furanyl, 2-thienyl, 3-thienyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 20 5-thiazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 4-pyrazolyl, 5-pyrazolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, oxa-2,4-diazolyl [sic], oxa-3,4-diazolyl [sic], thia-2,4-diazolyl [sic], thia-3,4-diazolyl [sic] and triazolyl, where the heteroaromatic rings can carry from one to 25 five halogen atoms as mentioned above, in particular fluorine and chlorine, and/or from one to three of the following radicals:

30 C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, cyano, nitro, C₁-C₈-alkylcarbonyl, C₁-C₈-alkoxycarbonyl, phenyl, phenoxy, phenylcarbonyl as mentioned above in general and in particular;

35 R⁴ is also phenyl or naphthyl which can be substituted by one or more, eg. from one to three, of the following radicals: halogen, nitro, cyano, hydroxyl, mercapto, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino, di-C₁-C₄-alkylamino, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, in particular as mentioned for 40 R⁷ and R⁸, and, for example, 3-hydroxyphenyl, 4-dimethylaminophenyl, 2-mercaptophenyl, 3-methoxycarbonylphenyl, 4-acetylphenyl, 1-naphthyl, 2-naphthyl, 3-bromo-2-naphthyl, 4-methyl-1-naphthyl, 5-methoxy-1-naphthyl, 6-trifluoromethyl-1-naphthyl, 7-chlor-1-naphthyl, 8-hydroxy-1-naphthyl; 45

or R⁴ and R⁵ form together with the adjacent carbon atom a 3- to 6-membered ring which can contain an oxygen or sulfur atom and is unsubstituted or carries from one to three, depending on the ring size, of the following radicals:

- 5 C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy, C₁-C₄-alkylthio as mentioned above in general and in particular;
- 10 R⁵ hydrogen, C₁-C₄-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl, C₃-C₈-cycloalkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxyalkyl, C₁-C₄-alkylthioalkyl or phenyl as mentioned above for R⁴ in particular;
- 15 R⁶ C₁-C₈-alkyl, C₃-C₆-alkenyl, C₃-C₆-alkynyl or C₃-C₈-cycloalkyl as mentioned above in particular, it being possible for each of these radicals to be substituted one or more times by: halogen, nitro, cyano, C₁-C₄-alkoxy, C₃-C₆-alkenyloxy, C₃-C₆-alkynyloxy, C₁-C₄-alkylthio, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl, C₁-C₄-alkoxycarbonyl, C₁-C₄-alkylamino, 20 di-C₁-C₄-alkylamino or unsubstituted or substituted phenyl or phenoxy as mentioned above in particular;
- 25 phenyl or naphthyl which can be substituted by one or more of the following radicals: halogen, nitro, cyano, hydroxyl, amino, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, phenoxy, C₁-C₄-alkylthio, C₁-C₄-alkylamino [sic] or C₁-C₄-dialkylamino as mentioned in particular for R⁷ and R⁴;
- 30 a five- or six-membered heteroaromatic ring which contains from one to three nitrogen atoms and/or one sulfur or oxygen atom and which can carry from one to four halogen atoms and/or one or two of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkyl- 35 thio, phenyl, phenoxy or phenylcarbonyl, where the phenyl radicals in turn can carry from one to five halogen atoms and/or from one to three of the following radicals: C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy and/or C₁-C₄-alkylthio as mentioned in particular for R⁴;
- 40 Y sulfur, oxygen or a single bond
- Z sulfur or oxygen.

45 Particularly preferred compounds of the formula I are those where R² and R³ are methoxy and X is CH. Also preferred are compounds of the formula I where R² and R³ are methoxy, X is CH, Y and Z are

oxygen and R^5 is C_1-C_4 -alkyl. The preferred radical in the case of R^1 is OR^{10} where R^{10} is hydrogen or C_1-C_4 -alkyl.

R^4 is particularly preferably C_1-C_4 -alkyl, unsubstituted or substituted phenyl or an aromatic heterocyclic radical containing one hetero atom, such as furyl or thienyl.

R^6 is particularly preferably phenyl which is unsubstituted or substituted 1-3 times by halogen, C_1-C_4 -alkyl, C_1-C_4 -alkoxy and/or C_1-C_4 -alkylthio.

Examples of preferred compounds are listed in the following Table.

Compounds 4.42 and 4.58 (Example 10, Tab. 4) are particularly preferably used according to the invention.

20

25

30

35

40

45

Table

R ¹	R ⁴	R ⁵	R ⁶	R ²	R ³	X	Y	Z
OH	Phenyl	Methyl	Methyl	OCH ₃	OCH ₃	CH	S	S
OH	Phenyl	Methyl	Methyl	OCH ₃	OCH ₃	CH	O	S
OCH ₃	Phenyl	Methyl	Methyl	OCH ₃	OCH ₃	CH	S	S
OH	Phenyl	i-Propyl	Methyl	OCH ₃	OCH ₃	CH	O	O
OCH ₃	2-Fluorophenyl	Ethyl	Methyl	OCH ₃	OCH ₃	CH	O	O
OC ₂ H ₅	3-Chlorophenyl	Propyl	Methyl	OCH ₃	OCH ₃	N	O	O
ON(CH ₃) ₂	4-Bromophenyl	i-Propyl	Methyl	CF ₃	CF ₃	CH	S	O
ON=C(CH ₃) ₂	2-Thienyl	Methyl	Methyl	OCF ₃	OCF ₃	CH	O	S
HNSO ₂ C ₆ H ₅	3-Thienyl	Methyl	Methyl	CH ₃	CH ₃	CH	O	O
NHPhenyl	2-Furyl	Methyl	Methyl	Cl	Cl	CH	O	O
ONa	3-Furyl	Methyl	Methyl	OCH ₃	-OCH ₂ -CH ₂ -		S	O
O-CH ₂ -C=CH	Phenyl	Ethyl	Ethyl	OCH ₃	CF ₃	CH	O	O
OH	Phenyl	Propyl	Propyl	OCH ₃	OCF ₃	CH	O	S
OCH ₃	Phenyl	i-Propyl	i-Propyl	OCH ₃	CH ₃	CH	O	O
OC ₂ H ₅	Phenyl	Methyl	s-Butyl	OCH ₃	Cl	CH	S	O
ON(CH ₃) ₂	2-Methylphenyl	Methyl	Methyl	OCH ₃	OCH ₃	CH	O	O
ON(CH ₃) ₂	3-Methoxyphenyl	Methyl	Methyl	OCH ₃	OCH ₃	CH	O	O
ON=C(CH ₃) ₂	4-Nitrophenyl	Methyl	Methyl	OCH ₃	OCH ₃	CH	O	O
NHPhenyl	2-Oxazolyl	Methyl	Methyl	CF ₃	CF ₃	N	S	O

R ¹	R ⁴	R ⁵	R ⁶	R ²	R ³	X	Y	Z
ONa	4-Oxazolyl	Methyl	3-Propenyl [sic]	OCF ₃	OCF ₃	N	O	S
O-CH ₂ -C≡CH	5-Oxazolyl	Methyl	3-Propynyl [sic]	CH ₃	CH ₃	N	O	O
OH	3-Isoxazolyl	Methyl	Cyclopentyl	Cl	Cl	N	O	O
OCH ₃	4-Isoxazolyl	Methyl	Cyclohexyl	OCH ₃	-O-CH ₂ -CH ₂ -		O	O
OC ₂ H ₅	5-Isoxazolyl	Methyl	Cyclopropylmethyl	OCH ₃	CF ₃	N	S	O
ON(CH ₃) ₂	Phenyl	Methyl	1-Phenyl-3-propynyl [sic]	OCH ₃	OCF ₃	N	O	S
ON=C(CH ₃) ₂	2-Hydroxyphenyl	Methyl	Methyl	OCH ₃	CH ₃	N	O	O
ONSO ₂ C ₆ H ₅	3-Trifluoromethylphenyl	Methyl	Methyl	OCH ₃	Cl	N	O	O
NHPhenyl	4-Dimethylaminophenyl	Methyl	Methyl	OCH ₃	OCH ₃	CH	S	O
ONa	2-Imidazolyl	Ethyl	Methyl	OCH ₃	OCH ₃	CH	S	S
O-CH ₂ -C≡CH	4-Imidazolyl	Propyl	Methyl	OCH ₃	OCH ₃	N	S	S
OH	3-Pyrazolyl	i-Propyl	Methyl	CF ₃	CF ₃	CH	O	S
OCH ₃	4-Pyrazolyl	Methyl	Methyl	OCF ₃	OCF ₃	CH	O	O
OC ₂ H ₅	Phenyl	Methyl	Trifluoroethyl	CH ₃	CH ₃	CH	O	O
ON(CH ₃) ₂	Phenyl	Methyl	Benzyl	Cl	Cl	CH	O	O
ON(CH ₃) ₂	Phenyl	Methyl	2-Methoxyethyl	OCH ₃	-O-CH ₂ -CH ₂ -		S	O
ON=C(CH ₃) ₂	Phenylpropyl	Methyl	3-Methoxycarbonyl- [sic]	OCH ₃	CF ₃	N	S	S
NH-Phenyl	2-Pyridyl	Methyl	2-Chloroethyl	OCH ₃	OCF ₃	N	S	S
ONa	3-Pyridyl	Methyl	Methyl	OCH ₃	CH ₃	N	O	O
O-CH ₂ -C≡CH	4-Pyridyl	Methyl	Methyl	OCH ₃	Cl	N	O	O

R ¹	R ⁴	R ⁵	R ⁶	R ²	R ³	X	Y	Z
OCH ₃	Phenyl	CH ₃	Phenyl	OCH ₃	OCH ₃	CH	O	O
OH	Phenyl	CH ₃	Phenyl	OCH ₃	OCH ₃	CH	O	O
OH	Phenyl	CH ₃	Phenyl	OCH ₃	-O-CH ₂ -CH ₂ -		O	O
OH	Phenyl	CH ₃	Phenyl	OCH ₃	OCH ₃	N	O	O
OH	Phenyl	CH ₃	Phenyl	OCH ₃	OCH ₃	CH	S	O
OH	Phenyl	CH ₃	Phenyl	OCH ₃	OCH ₃	CH	S	S
OH	Phenyl	CH ₃	Phenyl	OCH ₃	OCH ₃	CH	O	S
OH	Phenyl	H	Phenyl	OCH ₃	OCH ₃	CH	O	O
OH	Phenyl	i-Propyl	Phenyl	OCH ₃	OCH ₃	CH	O	O
OH	CH ₃	CH ₃	Phenyl	OCH ₃	OCH ₃	CH	O	O
OH		-(CH ₂) ₅ -	Phenyl	Phenyl	OCH ₃	CH	O	O
OH	Phenyl	CH ₃	2-Thiazolyl	OCH ₃	OCH ₃	CH	O	O
OH	2-Thienyl	CH ₃	Phenyl	OCH ₃	OCH ₃	CH	O	O
OCH ₃	2-Fluorophenyl	Ethyl	Phenyl	OCH ₃	OCH ₃	CH	O	O
OC ₂ H ₅	3-Chlorophenyl	Propyl	Phenyl	OCH ₃	OCH ₃	N	O	O
ON(CH ₃) ₂	4-Bromophenyl	i-Propyl	Phenyl	CF ₃	CF ₃	CH	S	O
ON=C(CH ₃) ₂	2-Thienyl	Methyl	Phenyl	OCF ₃	OCF ₃	CH	O	S
NH-SO ₂ -C ₆ H ₅	3-Thienyl	Methyl	Phenyl	CH ₃	CH ₃	CH	O	O
NHPhenyl	2-Furyl	Methyl	Phenyl	Cl	Cl	CH	O	O
ONa	3-Furyl	Methyl	Phenyl	OCH ₃	-O-CH ₂ -CH ₂ -		S	O
O-CH ₂ ≡CH	Phenyl	Ethyl	2-Fluorophenyl	OCH ₃	CF ₃	CH	O	O

R ¹	R ⁴	R ⁵	R ⁶	R ²	R ³	X	Y	Z
OH	Phenyl	Propyl	3-Chlorophenyl	OCH ₃	OCF ₃	CH	O	S
OCH ₃	Phenyl	i-Propyl	4-Bromophenyl	OCH ₃	CH ₃	CH	O	O
OC ₂ H ₅	Phenyl	Methyl	4-Thiazolyl	OCH ₃	Cl	CH	S	O
ON(CH ₃) ₂	2-Methylphenyl	Methyl	Phenyl	OCH ₃	OCH ₃	CH	O	O
ON=C(CH ₃) ₂	3-Methoxyphenyl	Methyl	Phenyl	OCH ₃	OCH ₃	CH	O	O
NH-SO-C ₆ H ₅	4-Nitrophenyl	Methyl	Phenyl	OCH ₃	OCH ₃	CH	O	O
NHPhenyl	Methyl	Methyl	Phenyl	CF ₃	CF ₃	N	S	O
ONa	Methyl	Methyl	2-Methylphenyl	OCF ₃	OCF ₃	N	O	S
O-CH ₂ -C≡CH	Methyl	Methyl	3-Methoxyphenyl	CH ₃	CH ₃	N	O	O
OH	Methyl	Methyl	4-Nitrophenyl	Cl	Cl	N	O	O
OCH ₃	Phenyl	Methyl	3-Imidazolyl	OCH ₃	-O-CH ₂ -CH ₂ -		O	O
OC ₂ H ₅	Phenyl	Methyl	4-Imidazolyl	OCH ₃	CF ₃	N	S	O
ON(CH ₃) ₂	Phenyl	Methyl	2-Pyrazolyl	OCH ₃	OCF ₃	N	O	S
ON=C(CH ₃) ₂	2-Hydroxyphenyl	Methyl	Phenyl	OCH ₃	CH ₃	N	O	O
NH-SO ₂ -C ₆ H ₅	3-Trifluoromethylphenyl	Methyl	Phenyl	OCH ₃	Cl	N	O	O
NHPhenyl	4-Dimethylaminophenyl	Methyl	Phenyl	OCH ₃	OCH ₃	CH	S	O
ONa	3-Imidazolyl	Ethyl	Phenyl	OCH ₃	OCH ₃	CH	S	S
O-CH ₂ -C≡CH	4-Imidazolyl	Propyl	Phenyl	OCH ₃	OCH ₃	N	S	S
OH	3-Pyrazolyl	i-Propyl	Phenyl	CF ₃	CF ₃	CH	O	S
OCH ₃	4-Pyrazolyl	Methyl	Phenyl	OCF ₃	OCF ₃	CH	O	O
OC ₂ H ₅	Phenyl	Methyl	2-Dimethylaminophenyl	CH ₃	CH ₃	CH	O	O

R ¹	R ⁴	R ⁵	R ⁶	R ²	R ³	X	Y	Z
ON(CH ₃) ₂	Phenyl	Methyl	3-Hydroxyphenyl	Cl	Cl	CH	O	O
ON=C(CH ₃) ₂	Phenyl	Methyl	4-Trifluoromethyl-phenyl	OCH ₃	-O-CH ₂ -CH ₂ -		S	O
NH-SO ₂ -C ₆ H ₅	Phenyl	Methyl	2-Oxazolyl	OCH ₃	CF ₃	N	S	S
NH-Phenyl	2-Pyridyl	Methyl	4-Isoxazolyl	OCH ₃	OCF ₃	N	S	S
ONa	3-Pyridyl	Methyl	Phenyl	OCH ₃	CH ₃	N	O	O
O-CH ₂ -C≡CH	4-Pyridyl	Methyl	Phenyl	OCH ₃	Cl	N	O	O

The compounds of the present invention provide a novel therapeutic potential for the treatment of hypertension, pulmonary hypertension, myocardial infarct, angina pectoris, acute kidney failure, renal insufficiency, cerebral vasospasms, cerebral ischemia, subarachnoid hemorrhage, migraine, asthma, atherosclerosis, endotoxic shock, endotoxin-induced organ failure, intravascular coagulation, restenosis after angioplasty and cyclosporin-induced kidney failure or hypertension.

10

The good effect of the compounds can be shown in the following experiments:

Receptor-binding studies

15

Cloned human ET_A receptor-expressing CHO cells and guinea pig cerebellar membranes with $> 60\%$ ET_B receptors compared with ET_A receptors were used for binding studies.

20 Membrane preparation

The ET_A receptor-expressing CHO cells were grown in F_{12} medium with 10% fetal calf serum, 1% glutamine, 100 U/ml penicillin and 0.2% streptomycin (Gibco BRL, Gaithersburg, MD, USA). After 48 h, the cells were washed with PBS and incubated with 0.05% trypsin-containing PBS for 5 min. The mixture was then neutralized with F_{12} medium and the cells were collected by centrifugation at 300 x g. For lysis of the cells, the pellet was briefly washed with lysis buffer (5 mM tris-HCl, pH 7.4 with 10% glycerol) and then incubated at a concentration of 10^7 cells/ml of lysis buffer at 4°C for 30 min. The membranes were centrifuged at 20,000 x g for 10 min, and the pellet was stored in liquid nitrogen.

35 Guinea pig cerebella were homogenized in a Potter-Elvehjem homogenizer and obtained by differential centrifugation at 1000 x g for 10 min and repeated centrifugation of the supernatant at 20,000 x g for 10 min.

40 Binding assays

For the ET_A and ET_B receptor binding assay, the membranes were suspended in incubation buffer (50 mM tris-HCl, pH 7.4 with 5 mM $MnCl_2$, 40 μ g/ml bacitracin and 0.2% BSA) at a concentration of 50 μ g of protein per assay mixture and incubated at 25°C with 25 pM [^{125}I]- ET_1 [sic] (ET_A receptor assay) or 25 pM [^{125}I]-RZ₃ [sic] (ET_B receptor assay) in the presence and absence of test substance. The non-specific binding was determined using

10⁻⁷ M ET₁. After 30 min, the free and the bound radioligand were separated by filtration through a GF/B glass fiber filter (Whatman, England) in a Skatron cell collector (Skatron, Lier, Norway), and the filters were washed with ice-cold tris-HCl buffer, 5 pH 7.4 with 0.2% BSA. The radioactivity collected on the filters was quantified using a Packard 2200 CA liquid scintillation counter.

The K_i values were determined by non-linear regression analysis 10 using the LIGAND program.

Table A shows the effect of compounds of the formula I as the K_i [mol/l] determined in the experiments.

T250X
15 Table A

Compound	K _i [mol/l]	
	ET-A	ET-B
4.42	2.5 × 10 ⁻⁷	3.0 × 10 ⁻⁶
4.58	1.6 × 10 ⁻⁷	4.7 × 10 ⁻⁶

Functional in vitro assay system for searching for endothelin re-
25 ceptor (subtype A) antagonists

This assay system is a functional cell-based assay for endothelin
30 receptors. Certain cells when stimulated with endothelin 1 (ET₁) show an increase in the intracellular calcium concentration. This increase can be measured in intact cells which have been loaded with calcium-sensitive dyes.

1-Fibroblasts which have been isolated from rats and in which an
35 endogenous endothelin receptor of subtype A has been detected were loaded with the fluorescent dye Fura 2-am as follows: after trypsinization the cells were resuspended in buffer A (120 mM NaCl, 5 mM KCl, 1.5 mM MgCl₂, 1 mM CaCl₂, 25 mM HEPES, 10 mM glucose, pH 7.4) to a density of 2 × 10⁶/ml and incubated with Fura 2-am (2 μM), Pluronic F-127 (0.04%) and DMSO (0.2%) at 37°C in
40 the dark for 30 min. The cells were then washed twice with buffer A and resuspended at 2 × 10⁶/ml.

The fluorescence signal at Ex/Em 380/510 from 2 × 10⁵ cells per ml
45 was recorded continuously at 30°C. The test substances were added to the cells and, after incubation with ET₁ for 3 min, the maximum change in the fluorescence was determined. The response of

the cells to ET1 without previous addition of a test substance served as control and was set equal to 100%.

Table B indicates the effect of the compounds of the formula I as 5 the IC₅₀ [mol/l] determined in the experiments.

T260X
Table B

10	Compound	IC ₅₀ [mol/l]
	4.42	7.4 * 10 ⁻⁷
	4.58	1.0 * 10 ⁻⁶

Testing of ET antagonists in vivo

15 Male SD rats weighing 250-300 g were anesthetized with amobarbital, artificially ventilated, vagotomized and pithed. The carotid artery and jugular vein were catheterized.

20 In control animals, intravenous administration of 1 µg/kg ET1 leads to a distinct rise in blood pressure which persists for a lengthy period.

25 5 min before administration of ET1, the test animals received the test compounds by i.v. injection (1 ml/kg). To determine the ET-antagonistic properties, the rise in blood pressure for the test animals was compared with that for the controls.

30 Endothelin-1-induced sudden death in mice

35 The test is based on the inhibition of the sudden heart death of mice which is caused by endothelin, probably by constriction of the coronary vessels, on pretreatment with endothelin receptor antagonists. Intravenous injection of 10 nmol/kg endothelin in a volume of 5 ml/kg of body weight is followed within a few minutes by the death of the animals.

40 The lethal endothelin-1 dose is checked in each case on a small group of animals. Intravenous administration of the test substance is followed, usually after 5 min, by the lethal endothelin-1 injection in the reference group. With other modes of administration the times between the doses are longer, where appropriate up to several hours.

45 The survival rate is recorded and effective doses for protection of 50% of the animals (ED 50) against endothelin-induced heart death for 24 h or longer are determined.

26

Functional vessel test for endothelin receptor antagonists

Initially, a contraction is induced by K^+ in segments of rabbit aorta after an initial tension of 2 g and a relaxation time of 5 1 h in Krebs-Henseleit solution at 37°C and pH 7.3 - 7.4. After washing out, an endothelin dose-response plot is constructed up to a maximum.

Potential endothelin antagonists are administered to other specimens of the same vessel 15 min before starting the endothelin dose-response plot. The effects of the endothelin are calculated as a % of the K^+ -induced contraction. Effective endothelin antagonists cause a shift to the right in the endothelin dose-response plot.

15

The compounds according to the invention can be administered orally or parenterally (subcutaneously, intravenously, intramuscularly, intraperitoneally) in a conventional way. Administration can also take place with vapors or sprays through the nasal pharyngeal space.

20

The dosage depends on the age, condition and weight of the patient and on the mode of administration. As a rule, the daily dose of active substance is about 0.5-50 mg/kg of body weight on 25 oral administration and about 0.1-10 mg/kg of body weight on parenteral administration.

The novel compounds can be administered in conventional solid or 30 liquid pharmaceutical forms, eg. uncoated or (film-)coated tablets, capsules, powders, granules, suppositories, solutions, ointments, creams or sprays. These are produced in a conventional way. For this purpose the active substances can be processed with conventional pharmaceutical aids such as tablet binders, fillers, 35 preservatives, tablet disintegrants, flow regulators, plasticizers, wetting agents, dispersants, emulsifiers, solvents, release-slowing agents, antioxidants and/or propellant gases (cf. H. Sucker et al.: Pharmazeutische Technologie, Thieme-Verlag, Stuttgart, 1991). The forms obtained in this way normally contain from 40 0.1 to 90% by weight of active substance.

~~Imp 83~~ Synthesis Examples

Synthesis of compounds of the general formula VI

5 Example 1

Methyl 3-methoxy-3-(3-methoxyphenyl)-2-hydroxybutyrate

19.5 g (88 mmol) of methyl 3-(3-methoxyphenyl)-2,3-epoxybutyrate are dissolved in 200 ml of absolute methanol, and 0.1 ml of boron
10 trifluoride etherate is added. The mixture is stirred at room temperature for 12 hours and the solvent is removed by distillation. The residue is taken up in ethyl acetate, washed with sodium bicarbonate solution and water and dried over sodium sulfate. After removal of the solvent by distillation, 21.1 g of a
15 pale yellow oil remain.

Yield: 94% (1:1 mixture of diastereomers)

Example 2

20 Methyl 3-benzyloxy-3-phenyl-2-hydroxybutyrate


9.6 g (50 mmol) of methyl 3-phenyl-2,3-epoxybutyrate are dissolved in 150 ml of benzyl alcohol, and 0.5 ml of concentrated sulfuric acid is added. The mixture is stirred at 50°C for 6 hours
25 and allowed to cool to room temperature. After neutralization with sodium bicarbonate solution, the excess benzyl alcohol is removed by distillation under high vacuum, and the residue is purified by flash chromatography on silica gel with 9:1 n-hexane/
30 ethyl acetate. After removal of the solvent by distillation, 6.5 g of a colorless oil remain.

Yield: 43% (3:2 mixture of diastereomers)

35 All the compounds mentioned in Table 1 were prepared in a similar way.

40

45

Table 1: Intermediates of the formula VI with R¹ = CH₃

	No.	R ⁶	R ⁴	R ⁵	DR*	M.p. [°C]
10	1.1	Methyl	3-Methoxyphenyl	Methyl	1:1	Oil
	1.2	Benzyl	Phenyl	Methyl	3:2	Oil
	1.3	Methyl	2-Fluorophenyl	Methyl	1:1	Oil
	1.4	Methyl	4-i-Propylphenyl	Methyl		
15	1.5	Methyl	2-Methylphenyl	Methyl	2:1	Oil
	1.6	Methyl	3-Methylphenyl	Methyl		
	1.7	Methyl	4-Methylphenyl	Methyl	3:2	Oil
	1.8	Methyl	3-Nitrophenyl	Methyl		
20	1.9	Methyl	4-Bromophenyl	Methyl	3:1	Oil
	1.10	Methyl	2-Furyl	Methyl		
	1.11	Methyl	3-Furyl	Methyl		
	1.12	Methyl	2-Thienyl	Methyl		
25	1.13	Methyl	3-Thienyl	Methyl		
	1.14	Methyl	2-Pyridyl	Methyl		
	1.15	Methyl	3-Pyridyl	Methyl		
	1.16	Methyl	4-Pyridyl	Methyl		
30	1.17	Methyl	2-Thiazolyl	Methyl		
	1.18	Methyl	3-Isoxazolyl	Methyl		
	1.19	Methyl	4-Imidazolyl	Methyl		
	1.20	Methyl	2-Pyrazolyl	Methyl		
35	1.21	Methyl	4-Chlorophenyl	Methyl	2:1	Oil
	1.22	Benzyl	3-Methylphenyl	Methyl	1:1	Oil
	1.23	Methyl	4-Fluorophenyl	Methyl	1:1	Oil
	1.24	Benzyl	4-Bromophenyl	Methyl	1:1	Oil
40	1.25	Benzyl	4-Chlorophenyl	Methyl	3:2	Oil
	1.26	Benzyl	4-Fluorophenyl	Methyl	1:1	Oil
	1.27	Methyl	Phenyl	Ethyl	1:1	Oil
	1.28	Methyl	3-Nitrophenyl	Methyl	2:1	Oil
45	1.29	Ethyl	4-Methylphenyl	Methyl	1:1	Oil
	1.30	Benzyl	4-Methylphenyl	Methyl	1:1	Oil
	1.31	Benzyl	Phenyl	Ethyl	1:0	Oil
	1.32	4-Fluorobenzyl	Phenyl	Methyl	1:1	Oil

* Diastereomer ratio

Synthesis of compounds of the general formula I:

Example 3:

Methyl 3-benzyloxy-3-phenyl-2-(4,6-dimethoxy-2-pyrimidinyl)oxy-
5 butyrate [sic]

3 g (10 mmol) of methyl 3-benzyloxy-3-phenyl-2-hydroxybutyrate
(comp. 1.1) are dissolved in 40 ml of dimethylformamide, and
0.3 g (12 mmol) of sodium hydride is added. The mixture is
10 stirred for 1 hour and then 2.2 g (10 mmol) of 4,6-dimethoxy-
2-methylsulfonylpyrimidine are added. The mixture is stirred at
room temperature for 24 hours and then cautiously hydrolyzed with
10 ml of water, the pH is adjusted to 5 with acetic acid, and the
solvent is removed by distillation under high vacuum. The residue
15 is taken up in 100 ml of ethyl acetate, washed with water, dried
over sodium sulfate and distilled to remove solvents. 10 ml of
methyl t-butyl ether are added to the residue, and the precipi-
tate is filtered off with suction. Drying results in 2.4 g of a
white powder.

20

Yield: 55% (1:1 mixture of diastereomers)

M.p.: 115 - 117°C

Example 4

25 3-Benzoyloxy-3-phenyl-2-(4,6-dimethoxy-2-pyrimidinyl)oxybutyric
[sic] acid

1.4 g (3 mmol) of methyl 3-benzyloxy-3-phenyl-2-(4,6-dimethoxy-
30 2-pyrimidinyl)oxybutyrate [sic] (Example 3) are dissolved in
20 ml of methanol and 20 ml of tetrahydrofuran, and 3.7 g of 10%
NaOH solution are added. The mixture is stirred at 60°C for 6
hours and at room temperature for 12 hours, the solvent is re-
moved by distillation under reduced pressure, and the residue is
35 taken up in 100 ml of water. The mixture is extracted with ethyl
acetate to remove unreacted ester. The aqueous phase is then ad-
justed to pH 1-2 with dilute hydrochloric acid and extracted with
ethyl acetate. After drying over magnesium sulfate and removal of
the solvent by distillation, a little acetone is added to the
40 residue, and the precipitate is filtered off with suction. Drying
results in 1.2 g of a white powder.

Yield: 88% (3:2 mixture of diastereomers)

M.p.: 165°C (decomposition)

45

Example 5

Methyl 3-benzyloxy-3-phenyl-2-[(4,6-dimethoxy-2-pyrimidinyl)thio]butyrate [sic]

5 11 g (25 mmol) of methyl 3-benzyloxy-3-phenyl-2-hydroxybutyrate (comp. 1.1) are dissolved in 50 ml of dichloromethane, 3 g (30 mmol) of triethylamine are added and, while stirring, 3.2 g (28 mmol) of methanesulfonyl chloride are added dropwise. The mixture is stirred at room temperature for 2 hours, washed with
10 water, dried over magnesium sulfate and concentrated under reduced pressure. The residue is taken up in DMF and added dropwise to a suspension of 12.9 g (75 mmol) of 4,6-dimethoxypyrimidine-2-thiol and 8.4 g (100 mmol) of sodium bicarbonate in 100 ml of DMF at 0°C. After stirring at room temperature for 2 hours and at
15 60°C for a further 2 hours, the mixture is poured into 1 l of ice-water, and the precipitate is filtered off with suction. Drying results in 3.2 g of a white powder.

Yield: 29% (1:1 mixture of diastereomers)

20

The compounds specified in Table 2 were prepared in a similar way to the above examples.

25

30

35

40

45




Table 2

No.	R ⁶	R ⁴	R ⁵	Y	R ¹	Diastereomers	M.p. (°C)
2.1	Benzyl	Phenyl	Methyl	O	OCH ₃	1:1	115-117
2.2	Benzyl	Phenyl	Methyl	O	OH	3:2	165 (decomp.)
2.3	Benzyl	Phenyl	Methyl	S	OCH ₃	1:1	
2.4	Benzyl	Phenyl	Methyl	S	OH		
2.5	Methyl	2-Fluorophenyl	Methyl	O	OCH ₃	1:1	126-128
2.6	Methyl	2-Fluorophenyl	Methyl	O	OH	2:1	185-186
2.7	Methyl	3-Methoxyphenyl	Methyl	O	OCH ₃	1:0 (5:1)	131-132 (93-95)
2.8	Methyl	3-Methoxyphenyl	Methyl	O	OH	1:0	187-189
2.9	Methyl	4-i-Propylphenyl	Methyl	O	OCH ₃		
2.10	Methyl	4-i-Propylphenyl	Methyl	O	OH		
2.11	Methyl	2-Methylphenyl	Methyl	O	OCH ₃	3:1	122-124
2.12	Methyl	2-Methylphenyl	Methyl	O	OH	1:1	135-137

No.	R ⁶	R ⁴	R ⁵	Y	R ¹	Diastereomers	M.p. (°C)
2.13	Methyl	3-Methylphenyl	Methyl	O	OCH ₃	1:1	105-110
2.14	Methyl	3-Methylphenyl	Methyl	O	OH	1:1	130-132
2.15	Methyl	4-Methylphenyl	Methyl	O	OCH ₃	1:1	99-102
2.16	Methyl	4-Methylphenyl	Methyl	O	OH	1:1	145-147
2.17	Methyl	4-Bromophenyl	Methyl	O	OCH ₃	1:0	148-150
2.18	Methyl	4-Bromophenyl	Methyl	O	OH	1:0	189-190
2.19	Methyl	2-Furyl	Methyl	O	OCH ₃		
2.20	Methyl	2-Furyl	Methyl	O	OH		
2.21	Methyl	3-Furyl	Methyl	O	OCH ₃		
2.22	Methyl	3-Furyl	Methyl	O	OH		
2.23	Methyl	2-Thienyl	Methyl	O	OCH ₃		
2.24	Methyl	2-Thienyl	Methyl	O	OH		
2.25	Methyl	2-Pyridyl	Methyl	O	OCH ₃	2:1	Oil
2.26	Methyl	2-Pyridyl	Methyl	O	ONa		175-176
2.27	Methyl	3-Pyridyl	Methyl	O	OCH ₃		
2.28	Methyl	3-Pyridyl	Methyl	O	OH		
2.29	Methyl	4-Pyridyl	Methyl	O	OCH ₃		
2.30	Methyl	4-Pyridyl	Methyl	O	OH		
2.31	Methyl	3-Chlorophenyl	Methyl	O	OCH ₃		
2.32	Methyl	3-Chlorophenyl	Methyl	O	OH		
2.33	Methyl	2-Thiazolyl	Methyl	O	OCH ₃		

No.	R ⁶	R ⁴	R ⁵	Y	R ¹	Diastereomers	M.p. (°C)
2.34	Methyl	2-Thiazolyl	Methyl	O	OH		
2.35	Methyl	3-Isoxazolyl	Methyl	O	OCH ₃		
2.36	Methyl	3-Isoxazolyl	Methyl	O	OH		
2.37	Methyl	4-Imidazolyl	Methyl	O	OCH ₃		
2.38	Methyl	4-Imidazolyl	Methyl	O	OH		
2.39	Methyl	2-Pyrazolyl	Methyl	O	OCH ₃		
2.40	Methyl	2-Pyrazolyl	Methyl	O	OH		
2.41	Benzyl	4-Chlorophenyl	Methyl	O	OCH ₃	1:1	112-114
2.42	Benzyl	4-Chlorophenyl	Methyl	O	OH		
2.43	i-Propyl	2-Fluorophenyl	Methyl	O	OCH ₃	4:1	115-120
2.44	i-Propyl	2-Fluorophenyl	Methyl	O	OH	2:1	143-145
2.45	Methyl	4-Fluorophenyl	Methyl	O	OCH ₃	1:1	122-125
2.46	Methyl	4-Fluorophenyl	Methyl	O	OH	3:1	170-172
2.47	Benzyl	3-Methylphenyl	Methyl	O	OCH ₃	1:1	94- 95
2.48	Benzyl	3-Methylphenyl	Methyl	O	OH	1:1	154-156
2.49	Methyl	4-Chlorophenyl	Methyl	O	OCH ₃	1:1	125-127
2.50	Methyl	4-Chlorophenyl	Methyl	O	OH	5:1	206-207
2.51	Methyl	Phenyl	Ethyl	O	OCH ₃	1:0	95-100
2.52	Methyl	Phenyl	Ethyl	O	OH	1:0	140-142
2.53	Benzyl	4-Fluorophenyl	Methyl	O	OCH ₃	1:1	95- 98
2.54	Benzyl	4-Fluorophenyl	Methyl	O	OH	4:1	153-154

No.	R ⁶	R ⁴	R ⁵	Y	R ¹	Diastereomers	M.p. (°C)
2.55	4-Fluoro-benzyl	Phenyl	Methyl	O	OCH ₃	1:0	152-153
2.56	4-Fluoro-benzyl	Phenyl	Methyl	O	OH	7:3	160-162
2.57	4-Bromobenzyl	Phenyl	Methyl	O	OCH ₃	9:1	158-160
2.58	4-Bromobenzyl	Phenyl	Methyl	O	OH	1:0	203-204
2.59	Benzyl	2-Fluorophenyl	Methyl	O	OCH ₃	1:0	129-130
2.60	Benzyl	2-Fluorophenyl	Methyl	O	OH	1:0	200-201
2.61	Benzyl	4-Bromophenyl	Methyl	O	OCH ₃	1:1	78- 79
2.62	Benzyl	4-Bromophenyl	Methyl	O	OH	1:1	156-158
2.63	Benzyl	4-Methylphenyl	Methyl	O	OCH ₃	1:1	Oil
2.64	Benzyl	4-Methylphenyl	Methyl	O	OH	4:1	158-159
2.65	Benzyl	Phenyl	Ethyl	O	OCH ₃	1:0	110-112
2.66	Benzyl	Phenyl	Ethyl	O	OH	1:0	92- 93
2.67	Ethyl	4-Methylphenyl	Methyl	O	OCH ₃	1:0	117-119
2.68	Ethyl	4-Methylphenyl	Methyl	O	OH	1:1	Oil
2.69	Methyl	2-Furyl	H	O	OCH ₃	1:1	Oil
2.70	Methyl	2-Furyl	H	O	OH	1:1	Oil
2.71	4-Chloro-benzyl	Phenyl	Methyl	O	OCH ₃	1:0	172-174
2.72	4-Chloro-benzyl	Phenyl	Methyl	O	OH	1:0	60- 61

No.	R ⁶	R ⁴	R ⁵	Y	R ¹	Diastereomers	M.p. (°C)
2.73	2-Butyl	4-Bromophenyl	Methyl	O	OCH ₃	-	104-106
2.74	2-Butyl	4-Bromophenyl	Methyl	O	OH	1:0	153-154
2.75	n-Propyl	4-Fluorophenyl	Methyl	O	OCH ₃	9:1	119-120
2.76	n-Propyl	4-Fluorophenyl	Methyl	O	OH	9:1	104-105
2.77	Methyl	3-Nitrophenyl	Methyl	O	OCH ₃	1:1	101-102
2.78	Methyl	3-Nitrophenyl	Methyl	O	OH	1:1	165-172
2.79	Methyl	4-Trifluorophenyl	Methyl	O	OCH ₃	1:0	112-113
2.80	Methyl	4-Trifluorophenyl	Methyl	O	OH	4:1	68- 70
2.81	Methyl	3-Thienyl	H	O	OCH ₃	1:1	80- 82
2.82	Methyl	3-Thienyl	H	O	OH	1:1	Oil
2.83	4-Chloro-benzyl	Phenyl	Methyl	O	OCH ₃	0:1	112-113
2.84	4-Chloro-benzyl	Phenyl	Methyl	O	OCH ₃	0:1	60- 61
2.85	Methyl	Phenyl	Ethyl	O	OCH ₃	1:3	125-130
2.86	Methyl	Phenyl	Ethyl	O	OH	0:1	133-135
2.87	Benzyl	3-Methoxyphenyl	Methyl	O	OCH ₃	3:1	86- 87
2.88	Benzyl	3-Methoxyphenyl	Methyl	O	OH	1:0	155
2.89	Benzyl	3-Methoxyphenyl	Methyl	O	OH	0:1	138-140
2.90	2-Phenylethyl	Phenyl	Methyl	O	OH	1:0	147-149
2.91	Methyl	3-Furyl	H	O	OCH ₃	1:1	Oil
2.92	Methyl	3-Furyl	H	O	OH	1:1	131-135

No.	R ⁶	R ⁴	R ⁵	Y	R ¹	Diastereomers	M.p. (°C)
2.93	3-CF ₃ -benzyl	Phenyl	Methyl	O	OCH ₃	2:1	151-152
2.94	3-CF ₃ -benzyl	Phenyl	Methyl	O	OH	1:1	Oil
2.95	2-Fluoro-benzene [sic]	Phenyl	Methyl	O	OCH ₃	2:1	170-173
2.96	2-Fluoro-benzene [sic]	Phenyl	Methyl	O	OH	1:0	160-162
2.97	2-Fluoro-benzyl	Phenyl	Methyl	O	OH	1:3	138-141
2.98	3-Fluoro-benzyl	Phenyl	Methyl	O	OCH ₃	1:1	81- 86
2.99	3-Fluoro-benzyl	Phenyl	Methyl	O	OH	4:1	195-197
2.100	3-Fluoro-benzyl	Phenyl	Methyl	O	ONa	3:1	250-260
2.101	4-Fluoro-benzyl	Phenyl	Methyl	O	OCH ₃	1:1	112-115
2.102	4-Fluoro-benzyl	Phenyl	Methyl	O	OH		

Synthesis of compounds of the general formula VI

Example 6

5

Methyl 3-phenoxy-3-phenyl-2-hydroxybutyrate

28.2 g (0.3 mol) of phenol and 19.2 g (0.1 mol) of methyl 3-phenyl-2,3-epoxybutyrate are heated together at 100°C for 6
10 hours. Removal of the excess phenol by distillation under high vacuum and purification of the residue by chromatography on silica gel with hexane/ethyl acetate mixtures result in 17.9 g of a pale yellow oil.

15 Yield: 62.5%

Example 7

Methyl 3-(4-bromophenyl)oxy-3-phenyl-2-hydroxybutyrate [sic]
20

51.9 g (0.3 mol) of 4-bromophenol and 19.2 g (0.1 mol) of methyl 3-phenyl-2,3-epoxybutyrate are stirred at 100°C for 8 h and at room temperature for 12 h. After removal of the excess phenol by distillation, the residue is purified by flash chromatography
25 (silica gel, n-hexane/ethyl acetate 9:1) to result in 7.2 g of a white solid.

Yield: 20%

M.p.: 133 - 135°C


30 The compounds specified in Table 3 were prepared in a similar way:

35

40

45

386

Table 3: Intermediates of the formula VI with R¹ = CH₃

		R ⁶	R ⁴	R ⁵	M.p. [°C]
10	3.1	Phenyl	Phenyl	Methyl	Oil
	3.2	4-Bromophenyl	Phenyl	Methyl	130–133
	3.3	Phenyl	Methyl	Methyl	
	3.4	Phenyl	Phenyl	i-Propyl	
15	3.5	2-Fluorophenyl	Phenyl	Methyl	
	3.6	3-Fluorophenyl	Phenyl	Methyl	Oil
	3.7	4-Fluorophenyl	Phenyl	Methyl	Oil
	3.8	4-Chlorophenyl	Phenyl	Methyl	
20	3.9	4-Nitrophenyl	Phenyl	Methyl	
	3.10	4-Methylphenyl	Phenyl	Methyl	Oil
	3.11	Phenyl	2-Fluorophenyl	Methyl	
	3.12	Phenyl	3-Methoxyphenyl	Methyl	
25	3.13	Phenyl	4-i-Propylphenyl	Methyl	
	3.14	Phenyl	2-Methylphenyl	Methyl	
	3.15	Phenyl	3-Nitrophenyl	Methyl	
	3.16	Phenyl	4-Bromophenyl	Methyl	
30	3.17	Phenyl	2-Furyl	Methyl	
	3.18	Phenyl	2-Thienyl	Methyl	Oil
	3.19	Phenyl	3-Furyl	Methyl	
	3.20	Phenyl	3-Thienyl	Methyl	
35	3.21	3-Methylphenyl	Phenyl	Methyl	Oil
	3.22	2-Methylphenyl	Phenyl	Methyl	Oil
	3.23	4-i-Propylphenyl	Phenyl	Methyl	Oil
	3.24	Phenyl	4-Chlorophenyl	Methyl	Oil

40

45

Synthesis of compounds of the general formula I:

Example 8

5 Methyl 3-phenoxy-3-phenyl-2-(4,6-dimethoxy-2-pyrimidinyl)oxybutyrate [sic]

4.4 g (15.4 mmol) of methyl 3-phenoxy-3-phenyl-2-hydroxybutyrate (compound 1.1) [sic] are dissolved in 40 ml of dimethylformamide, and 0.46 g (18.4 mmol) of sodium hydride is added. The mixture is stirred for 1 hour and then 3.4 g (15.4 mmol) of 4,6-dimethoxy-2-methylsulfonylpyrimidine are added. The mixture is stirred at room temperature for 24 hours and then cautiously hydrolyzed with 10 ml of water, the pH is adjusted to 5 with acetic acid, and the solvent is removed by distillation under high vacuum. The residue is taken up in 100 ml of ethyl acetate, washed with water, dried over sodium sulfate and distilled to remove solvents. 10 ml of methyl t-butyl ether are added to the residue, and the precipitate is filtered off with suction. Drying results in 1.6 g of a white powder.

Yield: 24.5%

M.p.: 143 - 145°C

25 Example 9

3-Phenoxy-3-phenyl-2-(4,6-dimethoxy-2-pyrimidinyl)oxybutyric [sic] acid

30 1.3 g of methyl 3-phenoxy-3-phenyl-2-(4,6-dimethoxy-2-pyrimidinyl)oxybutyrate [sic] (Example 8) are dissolved in 20 ml of MeOH and 40 ml of tetrahydrofuran, and 3.7 g of 10% NaOH solution are added. The mixture is stirred at 60°C for 6 hours and at room temperature for 12 hours, the solvent is removed by distillation under reduced pressure, and the residue is taken up in 100 ml of water. Unreacted ester is extracted with ethyl acetate. The aqueous phase is then adjusted to pH 1-2 with dilute hydrochloric acid and extracted with ethyl acetate. Drying over magnesium sulfate and removal of the solvent by distillation result in 40 1.0 g of a white powder.

Yield: 79.7%

M.p.: 50 - 55°C

Example 10


Methyl 3-phenoxy-3-phenyl-2-[(4,6-dimethoxy-2-pyrimidinyl)thio]butyrate [sic]

5

7.2 g (25 mmol) of methyl 3-phenoxy-3-phenyl-2-hydroxybutyrate (comp. 1.1) are dissolved in 50 ml of dichloromethane, 3 g (30 mmol) of triethylamine are added and, while stirring, 3.2 g (28 mmol) of methanesulfonyl chloride are added dropwise. The mixture is stirred at room temperature for 2 hours, washed with water, dried over magnesium sulfate and concentrated under reduced pressure. The residue is taken up in 100ml of DMF and added dropwise to a suspension of 12.9 g (75 mmol) of 4,6-dimethoxypyrimidine-2-thiol and 8.4 g (100 mmol) of sodium bicarbonate in 100 ml of DMF at 0°C. After stirring at room temperature for 2 hours and at 60°C for a further 2 hours, the mixture is poured into 1 l of ice-water, and the precipitate is filtered off with suction. Drying results in 4.2 g of a white powder.

20 Yield: 38%

The compounds specified in Table 4 were prepared in a similar way to the above examples.

T410X
25 Table 4

35	Ex. No.	R ⁶	R ⁴	R ⁵	R ¹	Y	M.p. [°C]
	4.1	Phenyl	Phenyl	Methyl	OCH ₃	O	100-103
	4.2	Phenyl	Phenyl	Methyl	OH	O	50-55
	4.3	Phenyl	Phenyl	Methyl	OCH ₃	S	
40	4.4	Phenyl	Phenyl	Methyl	OH	S	
	4.5	Phenyl	Phenyl	i-Propyl	OCH ₃	O	
	4.6	Phenyl	Phenyl	i-Propyl	OH	O	
	4.7	Phenyl	Methyl	Methyl	OCH ₃	O	
45	4.8	Phenyl	Methyl	Methyl	OH	O	
	4.9	4-Bromophenyl	Phenyl	Methyl	OCH ₃	O	130-135
	4.10	4-Bromophenyl	Phenyl	Methyl	OH	O	155-160

Ex. No.	R ⁶	R ⁴	R ⁵	R ¹	Y	M.p. [°C]
5	4.11	2-Fluorophenyl	Phenyl	Methyl	OCH ₃	O 128-134
	4.12	2-Fluorophenyl	Phenyl	Methyl	OH	O 170-171
	4.13	3-Fluorophenyl	Phenyl	Methyl	OCH ₃	O 85- 90
	4.14	3-Fluorophenyl	Phenyl	Methyl	OH	O 167-169
	4.15	4-Fluorophenyl	Phenyl	Methyl	OCH ₃	O 115-116
10	4.16	4-Fluorophenyl	Phenyl	Methyl	OH	O 122-125
	4.17	4-Chlorophenyl	Phenyl	Methyl	OCH ₃	O Oil
	4.18	4-Chlorophenyl	Phenyl	Methyl	OH	O 94- 98
	4.19	4-Methylphenyl	Phenyl	Methyl	OCH ₃	O 100-114
	4.20	4-Methylphenyl	Phenyl	Methyl	OH	O Oil
15	4.21	4-Nitrophenyl	Phenyl	Methyl	OCH ₃	O
	4.22	4-Nitrophenyl	Phenyl	Methyl	OH	O
	4.23	Phenyl	2-Fluorophenyl	Methyl	OCH ₃	O 130-132
	4.24	Phenyl	2-Fluorophenyl	Methyl	OH	O 194-195
	4.25	Phenyl	3-Methoxyphenyl	Methyl	OCH ₃	O Oil
20	4.26	Phenyl	3-Methoxyphenyl	Methyl	OH	O Oil
	4.27	Phenyl	4-i-Propylphenyl	Methyl	OCH ₃	O
	4.28	Phenyl	4-i-Propylphenyl	Methyl	OH	O
	4.29	Phenyl	4-Bromophenyl	Methyl	OCH ₃	O 129-131
	4.30	Phenyl	4-Bromophenyl	Methyl	OH	O Oil
25	4.31	Phenyl	2-Furyl	Methyl	OCH ₃	O
	4.32	Phenyl	2-Furyl	Methyl	OH	O
	4.33	Phenyl	3-Furyl	Methyl	OCH ₃	O
	4.34	Phenyl	3-Furyl	Methyl	OH	O
	4.35	Phenyl	2-Thienyl	Methyl	OCH ₃	O
30	4.36	Phenyl	2-Thienyl	Methyl	OH	O
	4.37	Phenyl	3-Thienyl	Methyl	OCH ₃	O
	4.38	Phenyl	3-Thienyl	Methyl	OH	O
	4.39	3-Methylphenyl	Phenyl	Methyl	OCH ₃	O 155
	4.40	3-Methylphenyl	Phenyl	Methyl	OH	O 100-101
35	4.41	4-i-Propyl-phenyl	Phenyl	Methyl	OCH ₃	O 130-131
	4.42	4-i-Propyl-phenyl	Phenyl	Methyl	OH	O 230
	4.43	Phenyl	4-Chlorophenyl	Methyl	OCH ₃	O 143-144
	4.44	Phenyl	4-Chlorophenyl	Methyl	OH	O 90- 92
	4.45	Phenyl	2-Methylphenyl	Methyl	OCH ₃	O 179-180
45	4.46	Phenyl	2-Methylphenyl	Methyl	OH	O
	4.47	2-Methylphenyl	Phenyl	Methyl	OCH ₃	O 95-114

Ex. No.	R ⁶	R ⁴	R ⁵	R ¹	Y	M.P. [°C]
5	4.48	2-Methylphenyl	Phenyl	Methyl	OH	O 80- 85
	4.49	Phenyl	4-Methylphenyl	Methyl	OCH ₃	O 110-112
	4.50	Phenyl	4-Methylphenyl	Methyl	OH	O 156-157
	4.51	Phenyl	3-Methylphenyl	Methyl	OCH ₃	O Oil
	4.52	Phenyl	3-Methylphenyl	Methyl	OH	O 158-160
10	4.53	4-Methoxy-phenyl	Phenyl	Methyl	OCH ₃	O 157-158
	4.54	4-Methoxy-phenyl	Phenyl	Methyl	OH	O 106-107
	4.55	Phenyl	4-Fluorophenyl	Methyl	OCH ₃	O 160-165
	4.56	Phenyl	4-Fluorophenyl	Methyl	OH	O 99-100
	4.57	4-Methylthio-phenyl	Phenyl	Methyl	OCH ₃	O 160-163
15	4.58	4-Methylthio-phenyl	Phenyl	Methyl	OH	O 248-250
	4.59	4-t-Butyl-phenyl	Phenyl	Methyl	OCH ₃	O 106-110
	4.60	4-t-Butyl-phenyl	Phenyl	Methyl	OH	O 250
	4.61	Phenyl	Phenyl	Ethyl	OCH ₃	O 115-117
	4.62	Phenyl	Phenyl	Ethyl	OH	O 84- 85
25	4.63	4-Acetoxy-phenyl	Phenyl	Methyl	OCH ₃	O 157-159
	4.64	4-Hydroxy-phenyl	Phenyl	Methyl	OH	O 80- 90

30

35

40

45